arXiv:1604.04682v1 [math.CA] 16 Apr 2016

A REPRESENTATION OF THE DICKSON POLYNOMIALS OF THE THIRD KIND BY LEGENDRE FUNCTIONS

NERANGA FERNANDO AND SOLOMON MANUKURE

Abstract. Let R be a commutative ring with identity. We show that the Dickson polynomials of the third kind $F_n(x,a)$ satisfy a non-homogeneous second order linear ordinary differential equation. We also show that the general solution to the differential equation reveals a connection between Dickson polynomials of the first and third kinds, and the Associated Legendre functions.

1. Introduction

Let n be a non-negative integer. The two elementary symmetric polynomials x_1+x_2 and x_1x_2 form a $\mathbb{Z}$-basis of the ring of symmetric polynomials in $\mathbb{Z}[x_1,x_2]$. In [2], it was shown that there exists $D_n(x,y)\in\mathbb{Z}[x,y]$ such that

$$x_1^n+x_2^n=D_n(x_1+x_2,x_1x_2).$$

The explicit form of $D_n(x,y)$ is given by the Waring's formula [3, Theorem 1.1]

$$D_n(x,y)=\sum_{i=0}^{\lfloor\frac{n}{2}\rfloor}\frac{n}{n-i}\binom{n-i}{i}(-y)^ix^{n-2i}.$$

Let q be a prime power and $\mathbb{F}_q$ denote the finite field with q elements. For fixed $a\in\mathbb{F}_q$, $D_n(x,a)\in\mathbb{F}_q[x]$ is the n-th Dickson polynomial of the first kind. Dickson polynomials have been widely studied for their permutation behavior over finite fields because of the applications of permutation polynomials in cryptography, coding theory, and combinatorics.

Let R be a commutative ring with identity and let $a\in R$. In [3], it is shown that the Dickson polynomials of the first kind, $D_n(x,a)$, satisfy the homogeneous second order ordinary differential equaion

$$(x^2-4a)D_n''(x,a)+xD_n'(x,a)-n^2\,D_n(x,a)=0.$$

The n-th Dickson polynomial of the second kind $E_n(x,a)$ is defined by

$$E_n(x,a)=\sum_{i=0}^{\lfloor\frac{n}{2}\rfloor}\binom{n-i}{i}(-a)^ix^{n-2i},$$

where $a\in R$ is a parameter. In [3], it is also shown that the Dickson polynomials of the second kind, $E_n(x,a)$, satisfy the homogeneous second order ordinary

2010 *Mathematics Subject Classification.* 34A30, 11T06.

Key words and phrases. Dickson polynomial, Differential Equation, Associated Legendre function, Hypergeometric function.

differential equaion

$$(x^2 - 4a)E_n''(x,a) + 3xE_n'(x,a) - n(n+2)\,E_n(x,a) = 0.$$

In [4], Stoll considered Dickson-type polynomials f_n over $\mathbb{R}$, which depend on two real parameters a and B. The explicit expression for f_n is given by

$$f_n(x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n + (B-2)i}{n-i} \binom{n-i}{i} (-a)^i\, x^{n-2i}. \tag{1.1}$$

We point out that (1.1) had also appeared earlier in [1].

Again in [4], Stoll showed that the Dickson-type polynomials f_n with $a \neq 0, B \in \mathbb{R}$ satisfy the second order homogeneous ordinary differential equation (See [4, Lemma 17])

$$(A_4x^4 + aA_2x^2 + a^2A_0)f_n'' + (B_3x^3 + aB_1x)f_n' - (C_2x^2 + aC_0)f_n = 0, \tag{1.2}$$

where $A_4, A_2, A_0, B_3, B_1, C_2, C_0 \in \mathbb{R}$.

For $a \in \mathbb{F}_q$, the n-th Dickson polynomial of the $(k+1)$-th kind $D_{n,k}(x,a)$ as defined by Wang and Yucas in [5] is

$$D_{n,k}(x,a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n-ki}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}. \tag{1.3}$$

Note that all the coefficients of $D_{n,k}(x,a)$ are integers and $D_{n,k}(x,a)$ also satisfies (1.2) with $A_4, A_2, A_0, B_3, B_1, C_2, C_0 \in \mathbb{Z}$.

In this paper, we consider the n-th Dickson polynomial of the third kind over a commutative ring R with identity. We thus hereafter assume that $a \in R$.

Our main goal is to explore the relationship between the Dickson polynomials of the third kind and associated Legendre functions. It turns out that the Dickson polynomials of the third kind satisfy the non-homogeneous second order linear ordinary differential equation given in Lemma 1.1. We show that the particular solution to the differential equation involves the Dickson polynomials of the first kind. The general solution to the associated homogeneous equation is enthralling since it involves the Associated Legendre functions which can be expressed in terms the gamma function and hypergeometric functions.

By (1.3), the n-th Dickson polynomial of the third kind $D_{n,2}(x,a)$ is given by

$$D_{n,2}(x,a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n-2i}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}, \tag{1.4}$$

where $a \in R$. Throughout the paper, we denote the n-th Dickson polynomial of the third kind $D_{n,2}(x,a)$ by $F_n(x,a)$.

The paper is organized as follows. In Section 2, we show that the Dickson polynomials of the third kind satisfy the non-homogeneous second-order linear ordinary differential equation given in the following lemma.

Lemma 1.1. *Let $a \neq 0$, $x = u + au^{-1}$ with $u \neq 0$ and $u^2 \neq a$. Then we have*

$$(x^2 - 4a)\; F_n''(x,a) + 3x\; F_n'(x,a) - n^2\; F_n(x,a) = 2n\; D_n(x,a), \tag{1.5}$$

where $D_n(x,a)$ is the n-th Dickson polynomial of the first kind.

In Section 3, we solve (1.5) and prove the following theorem.

Theorem 1.2. *Let $F_c(x,a)$ and $F_p(x,a)$ be the general solution to the associated homogeneous equation of* (1.5) *and the particular solution of* (1.5)*, respectively. Then*

$$F_c(x,a)=\frac{1}{\sqrt[4]{x^2-4a}}\Big[A\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)+B\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)\Big],$$

where A and B are constants, and $P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}$ and $Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}$ are the associated Legendre functions of the first and second kinds, respectively. Also,

$$F_p(x,a)=\sum_{k=0}^{n}b_k\,x^k,$$

where

$$\begin{cases} b_{k+2}=-\Big(\dfrac{n^2-k\,(k+2)}{4\,a\,(k+2)\,(k+1)}\Big)\,b_k, & n \text{ is odd}, k \text{ is even or } n \text{ is even}, k \text{ is odd},\\[2ex] b_{k+2}=\dfrac{1}{4\,a\,(k+2)\,(k+1)}\Big\{[k\,(k+2)-n^2]\,b_k-2n\,(\text{coefficient of}\,x^{n-2i}\,\text{in}\,D_n(x,a))\Big\}, & \\ \text{otherwise}, & \end{cases}$$

where $i=\frac{n-k}{2}$ and $D_n(x,a)$ is the n-th Dickson polynomial of the first kind.

In particular, we have

$$F_n(x,a)=F_c(x,a)+F_p(x,a).$$

1.1. **Preliminaries.** The Dickson polynomial $D_n(x,a)$ of the first kind of degree n in the indeterminate x and with parameter $a\in R$ is given as

$$D_n(x,a)=\sum_{i=0}^{\lfloor\frac{n}{2}\rfloor}\frac{n}{n-i}\binom{n-i}{i}(-a)^i x^{n-2i}.$$

Here $\lfloor\frac{n}{2}\rfloor$ denotes the largest integer $\le n/2$. Let u_1 and u_2 be indeterminates. Then Waring's formula (see [3, Theorem 1.1]) yields

$$u_1^n+u_2^n=\sum_{i=0}^{\lfloor\frac{n}{2}\rfloor}\frac{n}{n-i}\binom{n-i}{i}(-u_1u_2)^i(u_1+u_2)^{n-2i}$$

and thus

$$u_1^n+u_2^n=D_n(u_1+u_2,u_1u_2).$$

If we let $u_1=u$, $u_2=\frac{a}{u}$ and $x=u+\frac{a}{u}$, then the functional equation of the Dickson polynomial of the first kind is given by

$$D_n(u+\frac{a}{u},a)=u^n+\Big(\frac{a}{u}\Big)^n. \tag{1.6}$$

For $a \neq 0$, Let $x = u + au^{-1}$ with $u \neq 0$ and $u^2 \neq a$. Then the functional equation of the Dickson polynomial of the third kind, $F_n(x, a)$, is given by

$$F_n(x, a) = (u + \frac{a}{u}) \, \frac{u^n - \left(\frac{a}{u}\right)^n}{u - \frac{a}{u}}, \tag{1.7}$$

see [5, Eq. 4.1].

For m a real number, the Leibniz rule for the product of two functions $u(z)$ and $v(z)$ is given by

$$D_z^m uv \, = \, \sum_{\alpha=0}^{\infty} \binom{m}{\alpha} D_z^{m-\alpha} u \, D_z^{\alpha} v, \tag{1.8}$$

where

$$\binom{m}{\alpha} \, = \, \frac{\Gamma(m+1)}{\Gamma(m-\alpha+1)\,\Gamma(\alpha+1)}.$$

2. Proof of Lemma 1.1

From (1.7), we have

$$F_n(u + au^{-1}, a) \left(u - \frac{a}{u}\right) = (u + \frac{a}{u}) \left[u^n - \left(\frac{a}{u}\right)^n\right].$$

By differentiating both sides with respect to u, we get

$$\begin{aligned} &F_n'(u + au^{-1}, a) \left(u - \frac{a}{u}\right) \left(1 - \frac{a}{u^2}\right) + F_n(u + au^{-1}, a) \left(1 + \frac{a}{u^2}\right) \\ &= \left(u + \frac{a}{u}\right) \left[nu^{n-1} + \frac{na^n}{u^{n+1}}\right] + \left[u^n - \left(\frac{a}{u}\right)^n\right] \left(1 - \frac{a}{u^2}\right). \end{aligned}$$

Multiplying by u gives

$$\begin{aligned} &F_n'(u + au^{-1}, a) \left(u - \frac{a}{u}\right)^2 + F_n(u + au^{-1}, a) \left(u + \frac{a}{u}\right) \\ &= n \left(u + \frac{a}{u}\right) \left[u^n + \left(\frac{a}{u}\right)^n\right] + \left[u^n - \left(\frac{a}{u}\right)^n\right] \left(u - \frac{a}{u}\right). \end{aligned}$$

Differentiating both sides with respect to u again yields

$$\begin{aligned} &F_n''(u + au^{-1}, a) \left(u - \frac{a}{u}\right)^2 \left(1 - \frac{a}{u^2}\right) + 2 \, F_n'(u + au^{-1}, a) \left(u - \frac{a}{u}\right) \left(1 + \frac{a}{u^2}\right) \\ &+ F_n'(u + au^{-1}, a) \left(u + \frac{a}{u}\right) \left(1 - \frac{a}{u^2}\right) + F_n(u + au^{-1}, a) \left(1 - \frac{a}{u^2}\right) \\ &= n \left(u + \frac{a}{u}\right) \left[nu^{n-1} - \frac{na^n}{u^{n+1}}\right] + n \left[u^n + \left(\frac{a}{u}\right)^n\right] \left(1 - \frac{a}{u^2}\right) \\ &+ \left[u^n - \left(\frac{a}{u}\right)^n\right] \left(1 + \frac{a}{u^2}\right) + \left(u - \frac{a}{u}\right) \left[nu^{n-1} + \frac{na^n}{u^{n+1}}\right]. \end{aligned}$$

Multiplying by u gives

$$F_n''(u+au^{-1},a)\left(u-\frac{a}{u}\right)^3+3\ F_n'(u+au^{-1},a)\ \left[u^2-\left(\frac{a}{u}\right)^2\right]$$
$$+\ F_n(u+au^{-1},a)\ \left(u-\frac{a}{u}\right)$$
$$=(n^2+1)\ \left(u+\frac{a}{u}\right)\ \left[u^n-\left(\frac{a}{u}\right)^n\right]+2n\ \left(u-\frac{a}{u}\right)\ \left[u^n+\left(\frac{a}{u}\right)^n\right].$$

which implies

$$\left(u-\frac{a}{u}\right)^3\ F_n''(u+au^{-1},a)+3\ \left[u^2-\left(\frac{a}{u}\right)^2\right]\ F_n'(u+au^{-1},a)$$
$$+\left(u-\frac{a}{u}\right)\ F_n(u+au^{-1},a)$$
$$=(n^2+1)\ \left(u-\frac{a}{u}\right)\ F_n(u+au^{-1},a)+2n\ \left(u-\frac{a}{u}\right)\ \left[u^n+\left(\frac{a}{u}\right)^n\right].$$

Then we have

$$\left(u-\frac{a}{u}\right)^2\ F_n''(u+au^{-1},a)+3\ \left(u+\frac{a}{u}\right)\ F_n'(u+au^{-1},a)+F_n(u+au^{-1},a)$$
$$=(n^2+1)\ F_n(u+au^{-1},a)+2n\ \left[u^n+\left(\frac{a}{u}\right)^n\right].$$

which simplifies to

(2.1)
$$\left(u-\frac{a}{u}\right)^2\ F_n''(u+au^{-1},a)+3\ \left(u+\frac{a}{u}\right)\ F_n'(u+au^{-1},a)-n^2\ F_n(u+au^{-1},a)$$
$$=2n\ \left[u^n+\left(\frac{a}{u}\right)^n\right].$$

Note that, $D_n(x,a)=u^n+\left(\frac{a}{u}\right)^n$, where $D_n(x,a)$ is the n-th Dickson polynomial of the first kind (See (1.6)). Also note that, $\left(u-\frac{a}{u}\right)^2=x^2-4a$.

From 2.1 we have,

$$(x^2-4a)\ F_n''(x,a)+3x\ F_n'(x,a)-n^2\ F_n(x,a)=2n\ D_n(x,a).$$

Remark 2.1. *Let $a\neq 0$ and $x=u+au^{-1}$ with $u\neq 0$. When $u^2=a$, the functional equation of $F_n(a,x)$ is given by*

$$F_n(x,a)=2(\pm\sqrt{a})^n\,n; \tag{2.2}$$

see [5, Remark 2.4]. *In this case, we have $F_n'(x,a)=0$ which is not interesting.*

3. Proof of Theorem 1.2

In this section, we let $R=\mathbb{C}$.

In Section 2, we showed that Dickson polynomials of the third kind satisfy the second order non-homogeneous differential equation

$$(x^2-4a)\ F_n''(x,a)+3x\ F_n'(x,a)-n^2\ F_n(x,a)=2n\ D_n(x,a). \tag{3.1}$$

We first find the particular solution to (3.1), $F_p(x,a)$.

Let $F_p = \sum_{k=0}^{n} b_k\, x^k$ be the trial solution of the particular solution to (3.1). Then we have

$$F_p' = \sum_{k=1}^{n} k\, b_k\, x^{k-1} \qquad \text{and} \qquad F_p'' = \sum_{k=2}^{n} k(k-1)\, b_k\, x^{k-2}.$$

Then

(3.2)
$$\begin{aligned}
&(x^2-4a)\ F_n''(x,a) + 3x\ F_n'(x,a) - n^2\ F_n(x,a)\\
&= (x^2-4a)\ \sum_{k=2}^{n} k(k-1)\, b_k\, x^{k-2} + 3x\ \sum_{k=1}^{n} k\, b_k\, x^{k-1} - n^2\ \sum_{k=0}^{n} b_k\, x^k\\
&= \sum_{k=0}^{n-2} \left[k(k-1)\, b_k + 3\, k\, b_k - n^2\, b_k - 4a(k+2)(k+1)\, b_{k+2}\right] x^k + b_{n-1}x^{n-1} + 2nb_n x^n
\end{aligned}$$

Let $b_{n+1} = b_{n+2} = 0$. Now (3.2) can be written as

$$\begin{aligned}
&(x^2-4a)\ F_n''(x,a) + 3x\ F_n'(x,a) - n^2\ F_n(x,a)\\
&= \sum_{k=0}^{n} \left[k(k-1)\, b_k + 3\, k\, b_k - n^2\, b_k - 4a(k+2)(k+1)\, b_{k+2}\right] x^k
\end{aligned}$$

From (3.1) we have

$$\begin{aligned}
&\sum_{k=0}^{n} \left[k(k-1)\, b_k + 3\, k\, b_k - n^2\, b_k - 4a(k+2)(k+1)\, b_{k+2}\right] x^k\\
&= 2n \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}.
\end{aligned} \tag{3.3}$$

Consider the following cases.

(i) n is odd and k is even,
(ii) n is even and k is odd,
(iii) n is odd and k is odd, and
(iv) n is even and k is even.

In cases (i) and (ii), by comparing the coefficients in (3.3) we have

$$k(k-1)\, b_k + 3\, k\, b_k - n^2\, b_k - 4a(k+2)(k+1)\, b_{k+2} = 0$$

which implies

$$b_{k+2} = -\Big(\frac{n^2 - k\,(k+2)}{4\,a\,(k+2)\,(k+1)}\Big)\, b_k.$$

Now consider cases (iii) and (iv). By comparing the coefficients in (3.3) we have

$$b_{k+2} = \frac{1}{4\,a\,(k+2)\,(k+1)} \Big\{ \left[k\,(k+2) - n^2\right] b_k - 2n\,(\text{coefficient of}\, x^{n-2i}\, \text{in}\, D_n(x,a)) \Big\},$$

where $i = \frac{n-k}{2}$.

Now let's consider the associated homogeneous equation, i.e.

$$(x^2-4a)\ F_n''(x,a)+3x\ F_n'(x,a)-n^2\ F_n(x,a)=0. \tag{3.4}$$

Denote $F_n(x,a)$ by $F(x)$. Then

$$\begin{gathered}(x^2-4a)\ F''(x)+3x\ F'(x)-n^2\ F(x)=0\\ \frac{d}{dx}[(x^2-4a)^{\frac{3}{2}}\ F'(x)]-n^2\,(x^2-4a)^{\frac{1}{2}}\ F(x)=0\\ (4a)^{3/2}\frac{d}{dx}\Big[\Big(\frac{x^2}{4a}-1\Big)^{\frac{3}{2}}\ F'(x)\Big]-(4a)^{1/2}n^2\ \Big(\frac{x^2}{4a}-1\Big)^{\frac{1}{2}}\ F(x)=0\\ 4a\frac{d}{dx}\Big[\Big(\Big(\frac{x}{2\sqrt{a}}\Big)^2-1\Big)^{\frac{3}{2}}\ F'(x)\Big]-n^2\ \Big(\Big(\frac{x}{2\sqrt{a}}\Big)^2-1\Big)^{\frac{1}{2}}\ F(x)=0\end{gathered}$$

Let $X=\dfrac{x}{2\sqrt{a}}$. Then $dx=2\sqrt{a}\,dX$. It follows that

$$\begin{gathered}(4a)\frac{1}{2\sqrt{a}}\frac{d}{dX}\Big[\Big(X^2-1\Big)^{\frac{3}{2}}\ F'(2\sqrt{a}X)\Big]-n^2\ \Big(X^2-1\Big)^{\frac{1}{2}}\ F(2\sqrt{a}X)=0\\ 2\sqrt{a}\frac{d}{dX}\Big[\Big(X^2-1\Big)^{\frac{3}{2}}\ F'(2\sqrt{a}X)\Big]-n^2\ \Big(X^2-1\Big)^{\frac{1}{2}}\ F(2\sqrt{a}X)=0\end{gathered}$$

Let $P(X)=F(2\sqrt{a}X)$.

Then $P'(X)=2\sqrt{a}F'(2\sqrt{a}X)$ which implies $\dfrac{1}{2\sqrt{a}}P'(X)=F'(2\sqrt{a}X)$. Then we have

$$\frac{d}{dX}\Big[\Big(X^2-1\Big)^{\frac{3}{2}}\ P'(X)\Big]-n^2\ \Big(X^2-1\Big)^{\frac{1}{2}}\ P(X)=0. \tag{3.5}$$

Now consider the Legendre equation:

$$(1-x^2)P_l''(x)-2x\,P_l'(x)+l(l+1)P_l(x)=0$$

or equivalently

$$(x^2-1)P_l''(x)+2x\,P_l'(x)-l(l+1)P_l(x)=0. \tag{3.6}$$

Differentiating m times using Leibniz rule (see (1.8)) gives the following.

$$\begin{aligned}D_x^m[(x^2-1)\,P_l''(x)] &= \sum_{\alpha=0}^{\infty}\frac{\Gamma(m+1)}{\Gamma(m-\alpha+1)\Gamma(\alpha+1)}D_x^{\alpha}(x^2-1)D^{m-\alpha}P_l''(x)\\ &= (x^2-1)P_l^{(m+2)}(x)+\frac{\Gamma(m+1)}{\Gamma(m)\Gamma(2)}2x\ P_l^{(m+1)}(x)+\frac{\Gamma(m+1)}{\Gamma(m-1)\Gamma(3)}2\ P_l^{(m)}(x)\\ &= (x^2-1)P_l^{(m+2)}(x)+2mx\ P_l^{(m+1)}(x)+m(m-1)\ P_l^{(m)}(x).\end{aligned} \tag{3.7}$$

Similarly,

$$D_x^m[2x\,P_l'(x)]=2x\,P_l^{(m+1)}(x)+2m\,P_l^{(m)}(x). \tag{3.8}$$

(3.6), (3.7), and (3.8) yield

$$(x^2-1)P_l^{(m+2)}(x)+2m\,x\;P_l^{(m+1)}(x)+m(m-1)\;P_l^{(m)}(x)+2x\,P_l^{(m+1)}(x)$$
$$+2m\,P_l^{(m)}(x)-l(l+1)P_l^{(m)}(x)=0.$$

$$(x^2-1)P_l^{(m+2)}(x)+(2m+2)\,x\;P_l^{(m+1)}(x)-[l(l+1)-m(m+1)]P_l^{(m)}(x)=0.$$

$$(x^2-1)P_l^{(m+2)}(x)+2(m+1)\,x\;P_l^{(m+1)}(x)-(l-m)(l+m+1)P_l^{(m)}(x)=0. \tag{3.9}$$

Now, let $y(x)=P_l^{(m)}$. Then (3.9) becomes

$$(x^2-1)\frac{d^2y}{dx^2}+2(m+1)\,x\;\frac{dy}{dx}-(l-m)(l+m+1)\,y=0. \tag{3.10}$$

By multiplying (3.10) by $(x^2-1)^m$, we get

$$(x^2-1)^{m+1}\frac{d^2y}{dx^2}+2(m+1)\,x\,(x^2-1)^m\;\frac{dy}{dx}-(l-m)(l+m+1)\,(x^2-1)^m\,y=0$$

which can be rewritten in the form

$$\frac{d}{dx}\Big[\Big(x^2-1\Big)^{m+1}\;\frac{dy}{dx}\Big]-(l-m)(l+m+1)\,\Big(x^2-1\Big)^m\,y=0. \tag{3.11}$$

Now, let

$$y(x)=(x^2-1)^{-\frac{m}{2}}\,u(x). \tag{3.12}$$

Then

$$\frac{dy}{dx}=(x^2-1)^{-\frac{m}{2}}\;\frac{dy}{dx}-mx(x^2-1)^{-\frac{m}{2}-1}\,u.$$

Thus (3.11) becomes

$$\frac{d}{dx}\Big[\Big(x^2-1\Big)^{\frac{m}{2}+1}\;\frac{du}{dx}-mx\,(x^2-1)^{\frac{m}{2}}\,u\Big]-(l-m)(l+m+1)\,\Big(x^2-1\Big)^{\frac{m}{2}}\,u=0.$$

$$\Big(x^2-1\Big)^{\frac{m}{2}+1}\;\frac{d^2u}{dx^2}+\Big(\frac{m}{2}+1\Big)(2x)\Big(x^2-1\Big)^{\frac{m}{2}}\;\frac{du}{dx}-mx\Big(x^2-1\Big)^{\frac{m}{2}}\;\frac{du}{dx}-m\Big(x^2-1\Big)^{\frac{m}{2}}\,u$$
$$-mx\,\frac{m}{2}\,(2x)\,\Big(x^2-1\Big)^{\frac{m}{2}-1}\,u-(l-m)(l+m+1)\,\Big(x^2-1\Big)^{\frac{m}{2}}\,u=0.$$

Dividing by $\Big(x^2-1\Big)^{\frac{m}{2}}$, we obtain

$$(x^2-1)\,\frac{d^2u}{dx^2}+(m+2)x\,\frac{du}{dx}-mx\,\frac{du}{dx}-mu-m^2\frac{x^2}{x^2-1}\,u-(l-m)(l+m+1)\,u=0.$$

$$(x^2-1)\,\frac{d^2u}{dx^2}+2x\,\frac{du}{dx}-mu-m^2\Big(\frac{x^2-1+1}{x^2-1}\Big)\,u-(l-m)(l+m+1)\,u=0.$$

$$(x^2-1)\,\frac{d^2u}{dx^2}+2x\,\frac{du}{dx}-mu-m^2\Big(1+\frac{1}{x^2-1}\Big)\,u-(l-m)(l+m+1)\,u=0.$$

$$(x^2-1)\,\frac{d^2u}{dx^2}+2x\,\frac{du}{dx}-\Big[(l-m)(l+m+1)+m+m^2+\frac{m^2}{x^2-1}\Big]\,u=0.$$

$$(x^2-1)\,\frac{d^2u}{dx^2}+2x\,\frac{du}{dx}-\Big[l(l+1)+\frac{m^2}{x^2-1}\Big]\,u=0. \tag{3.13}$$

Eq. (3.13) is the so-called associated Legendre differential equation, with $0 \le m \le l$. It's general solution is given by (see [6, Ch. 5])

$$u_g = c_1\,P_l^{(m)}(x)\;+\;c_2\,Q_l^{(m)}(x),$$

where c_1 and c_2 are arbitrary constants, and $P_l^{(m)}(x)$ and $Q_l^{(m)}(x)$ are the associated Legendre functions of the first and second kinds, respectively. Thus, according to (3.12)

$$y_g(x)=\Big(x^2-1\Big)^{-\frac{m}{2}}\,u_g(x)$$

is the general solution to (3.11).

Now consider the equation

$$\frac{d}{dx}\Big[\Big(x^2-1\Big)^{\frac{3}{2}}\,\frac{dy}{dx}\Big]-n^2\,\Big(x^2-1\Big)^{\frac{1}{2}}\,y=0. \tag{3.14}$$

This is in the form of (3.5) with $m=\frac{1}{2}$ and $(l-m)(l+m+1)=n^2$. Since $l\ge 0$, we have $l=\sqrt{n^2+1}-\frac{1}{2}$. Then the solution to (3.14) is

$$y=(x^2-1)^{-\frac{1}{4}}\,\Big[c_1\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(x)+c_2\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(x)\Big].$$

So the solution to (3.5) is

$$P(X)=(X^2-1)^{-\frac{1}{4}}\,\Big[c_1\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(X)+c_2\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(X)\Big].$$

Since $X=\dfrac{x}{2\sqrt{a}}$ and $P(X)=F(2\sqrt{a}X)$, we have

$$\begin{aligned}F(x)&=\Big(\frac{x^2}{4a}-1\Big)^{-\frac{1}{4}}\,\Big[c_1\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)+c_2\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)\Big]\\&=(x^2-4a)^{-\frac{1}{4}}\,\Big[\frac{c_1}{(4a)^{-\frac{1}{4}}}\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)+\frac{c_2}{(4a)^{-\frac{1}{4}}}\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)\Big].\end{aligned}$$

If we denote the general solution to (3.4) by $F_c(x,a)$, then we have

$$F_c(x,a)=\frac{1}{\sqrt[4]{x^2-4a}}\,\Big[A\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)+B\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)\Big]$$

where A and B are arbitrary constants. Hence the general solution to the differential equation

$$(x^2-4a)\;F_n''(x,a)+3x\;F_n'(x,a)-n^2\;F_n(x,a)=2n\;D_n(x,a),$$

is

$$F_n(x,a)=F_c(x,a)+F_p(x,a),$$

i.e.

(3.15)

$$F_n(x,a) = \frac{1}{\sqrt[4]{x^2-4a}}\left[A\,P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big) + B\,Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}\Big(\frac{x}{2\sqrt{a}}\Big)\right] + \sum_{k=0}^{n} b_k\,x^k,$$

where b_k is determined as explained in Theorem 1.2.

In terms of hypergeometric functions, the Legendre functions can be written as (see [6, Ch. 5])

(3.16)

$$P^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(z) = \frac{1}{\Gamma(1-\mu)}\,\Big[\frac{1+z}{1-z}\Big]^{\frac{1}{4}}\,{}_2F_1\Big(-\sqrt{n^2+1}+\frac{1}{2},\sqrt{n^2+1}+\frac{1}{2};\frac{1}{2};\frac{1-z}{2}\Big),$$

for $\mid 1-z \mid < 2$, and

(3.17)

$$\begin{aligned} &Q^{\frac{1}{2}}_{\sqrt{n^2+1}-\frac{1}{2}}(z) \\ &= \frac{\sqrt{\pi}\,\Gamma(\sqrt{n^2+1}+1)}{2^{\sqrt{n^2+1}+\frac{1}{2}}\,\Gamma(\sqrt{n^2+1}+1)}\;\frac{e^{\frac{i\pi}{2}}(z^2-1)^{\frac{1}{4}}}{z^{\sqrt{n^2+1}+1}}\;{}_2F_1\left(\frac{\sqrt{n^2+1}+1}{2},\frac{\sqrt{n^2+1}+2}{2};\sqrt{n^2+1}+1;\frac{1}{z^2}\right) \\ &= \frac{\sqrt{\pi}}{2^{\sqrt{n^2+1}+\frac{1}{2}}}\;\frac{i\,(z^2-1)^{\frac{1}{4}}}{z^{\sqrt{n^2+1}+1}}\;{}_2F_1\left(\frac{\sqrt{n^2+1}+1}{2},\frac{\sqrt{n^2+1}+2}{2};\sqrt{n^2+1}+1;\frac{1}{z^2}\right) \end{aligned}$$

for $\mid z \mid > 1$,

where Γ denotes the gamma function and ${}_2F_1(a,b;c;z)$ is the hypergeometric function defined by the power series

$${}_2F_1(a,b;c;z) = \sum_{n=0}^{\infty}\frac{(a)_n\,(b)_n}{(c)_n}\frac{z^n}{n!},$$

for $\mid z \mid < 1$.

Here $(\cdot)_n$ denotes the rising factorial defined by

$$(a)_n = \begin{cases} 1 & , \quad n=0 \\ a(a+1)\cdots(a+n-1) & , \quad n>0. \end{cases}$$

Equations (3.15) - (3.17) determine the constants A and B.

References

[1] A. Dujella, I. Gusić, *Decomposition of a recursive family of polynomials*, Monatsh. Math. **152** (2007), no. 2, 97 – 104.

[2] I. G. Macdonald, *Symmetric Functions and Orthogonal Polynomials*, American Mathematical Society, Providence, RI, 1998.

[3] R. Lidl, G. L. Mullen, G. Turnwald, *Dickson polynomials*, Longman Scientific and Technical, Essex, United Kingdom, 1993.

[4] T. Stoll, *Complete decomposition of Dickson-type polynomials and related Diophantine equations*, Journal of Number Theory **128** (2008), 1157 – 1181.

[5] Q. Wang, J. L. Yucas, *Dickson polynomials over finite fields*, Finite Fields Appl. **18** (2012), 814 – 831.

[6] Z. X. Wang, D. R. Guo, *Special Functions*, World Scientific Publishing Co. Pte. Ltd., 1989.

[7] G. M. Wysin, *Associated Legendre Functions and Dipole Transition Matrix Elements*, online lecture notes.

Department of Mathematics, Northeastern University, Boston, MA 02115
E-mail address: w.fernando@neu.edu

Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620
E-mail address: smanukure@mail.usf.edu